

## Case Report

# Functionalization of a Volume-Stable Collagen Matrix Using Liquid Platelet-Rich Fibrin: A Case Report Presenting a New Approach for Root Coverage

**Roberta Michels , Gabriel Leonardo Magrin , ACC Cruz , RS Magini ,  
and Cesar Augusto Magalhães Benfatti **

*Center for Education and Research on Dental Implants (CEPID), Post-Graduate Program in Dentistry (PPGO), Federal University of Santa Catarina (UFSC), Florianópolis, Brazil*

Correspondence should be addressed to Gabriel Leonardo Magrin; [gabriel.magrin@posgrad.ufsc.br](mailto:gabriel.magrin@posgrad.ufsc.br)

Received 11 October 2022; Revised 29 January 2023; Accepted 4 February 2023; Published 22 March 2023

Academic Editor: Jamil Awad Shibli

Copyright © 2023 Roberta Michels et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This case report presents a novel approach for root coverage of multiple gingival recessions with a volume-stable collagen matrix functionalized with injectable platelet-rich fibrin (i-PRF). A patient with multiple gingival recessions in the anterior maxilla was submitted to root coverage by coronally advanced flap with split–full–split incisions. Blood collection was performed before surgery and i-PRF was obtained after centrifugation (relative centrifugal force (RCF) 400 g, 2700 rpm, 3 minutes). A volume-stable collagen matrix was soaked with i-PRF and applied as a substitute for autogenous connective tissue graft. A mean root coverage of 83% was observed after a 12-month follow-up period, and only slight modifications were detected in a 30-month follow-up consultation. The association of a volume-stable collagen matrix with i-PRF successfully treated multiple gingival recessions with reduced morbidity since a connective tissue collection was avoided.

## 1. Introduction

Gingival recession is a condition that can be defined as the apical migration of gingival tissue from the cemento-enamel junction and is associated with the loss of supportive periodontium [1]. Consequently, root exposure to the oral environment is observed, and patients can present dentin hypersensitivity, root caries, and aesthetics complaints [2]. Gingival recession can be caused by several factors like traumatic toothbrushing, inadequate tooth position (i.e., crowded teeth), orthodontic movement, deep cervical restorative margins, and persistent gingival inflammation [3]. Root coverage techniques have been employed to treat gingival recessions. According to the interproximal involvement, this condition can be classified as: RT1—no interproximal insertion loss; RT2—interproximal insertion loss not higher than the buccal level; and RT3—interproximal insertion loss higher than the buccal level [4]. The best prognosis has been observed in RT1 lesions [4, 5].

Coronally advanced flap (CAF) associated with autogenous connective tissue graft has demonstrated long-term effectiveness in multiple gingival recession treatment [6]. However, autogenous graft collection has limitations such as the risk of hemorrhage, limited availability of donor tissue, anatomical variations between patients, and postoperative morbidity [7]. Taking those points into consideration, alternative biomaterials have been developed for root coverage with stable outcomes and less invasive techniques, such as acellular dermal matrix, xenogeneic collagen matrix, enamel matrix derivative, and platelet aggregates [8, 9]. Xenogeneic collagen matrix has been used as a substitute for autogenous connective tissue graft; however, while autogenous tissue has viable fibroblasts and signaling molecules for stable long-term outcomes when associated with root coverage procedures [10, 11], xenogeneic matrices only mimic the extracellular matrix and do not have a robust evidence in terms of root coverage results [12]. In this context, adding signaling molecules into collagen matrices may increase cell migration,




(a)



(b)



(c)



(d)



(e)

FIGURE 1: Continued.



FIGURE 1: Preoperative evaluation and surgical procedure. (a) Clinical view before surgery, (b) a surgical flap with a split–full–split thickness configuration was raised, (c) volume-stable collagen matrix (Fibro-Gide, Geistlich Pharma AG, Wolhusen, Switzerland) before platelet-rich fibrin functionalization, (d) functionalization of the collagen matrix with injectable platelet-rich fibrin (i-PRF), (e) collagen matrix after functionalization, (f) positioning of the matrix impregnated with injectable platelet-rich fibrin (i-PRF) in the surgical site, (g) tension-free flap repositioning and sutures.

adhesion, and differentiation, improving patient healing and the overall root coverage result, as suggested in observations of a proof-of-concept study with acellular dermal matrix allografts [13].

Platelet-rich fibrin (PRF) is a blood-derived reservoir of bioactive molecules that are known to promote wound healing and tissue regeneration [14]. This platelet concentrate can act as a delivery system of molecules such as transforming growth factor (TGF)- $\beta$  and collagen type 1, which increases the release of other growth factors and signaling proteins [15]. Moreover, the PRF has been extensively studied in periodontology due to the potential for keratinization in surgical sites [16, 17]. Studies reported that injectable PRF (i-PRF) stimulates more vascularization and cell chemotaxis than other PRF preparations [15, 18]. Therefore, the aim of this case report was to describe a multiple gingival recession case treated by CAF associated with a volume-stable collagen matrix functionalized with i-PRF.

## 2. Case Presentation

**2.1. Baseline Examination.** A 26-year-old male patient sought the Center for Education and Research on Dental Implants at the Federal University of Santa Catarina (CEPID/UFSC) in February 2020, due to multiple gingival recessions in the aesthetic area of the maxilla caused by orthodontic treatment and brushing trauma. During clinical

examination, a thin gingival phenotype was detected (Figure 1(a)). Since no interproximal involvement was present, the gingival recessions were classified as RT1 [4]. Probing depths were  $\leq 3$  mm for all periodontal sites, bleeding on probing was present in 27.78% of the sites, and plaque index was 15%. The patient was diagnosed with biofilm-induced gingivitis [19].

**2.2. Ethical Aspects.** The patient was informed about the characteristics of the intervention and the data collection both orally and through a written information sheet. An informed consent was signed by the patient. No institutional approval of the ethics committee was necessary for this case report.

**2.3. Preoperative Clinical Sessions.** Before the surgical procedure, gingival health was obtained. The patient received brushing technique instructions, and gingivitis was treated with non-surgical periodontal therapy involving ultrasonic debridement, root planning, and professional prophylaxis. The patient was eligible for root coverage intervention after gingivitis treatment.

**2.4. Centrifugation Protocol for I-PRF.** Protocol for the i-PRF acquisition was performed before surgery, as previously described [15]. Briefly, in the surgical room, a nurse collected the blood from the patient in two plastic tubes that were immediately placed in a centrifuge (Intra-Lock L-PRF





FIGURE 2: Postoperative follow-up. (a) Clinical view on 2-month follow-up, (b) clinical observation on 9-month follow-up, (c) clinical view on 12-month follow-up.

Intra Spin, Birmingham, AL, USA) at 2700 rpm and 400 g relative centrifugal force (RCF) for 3 minutes. The yellow portion, the fibrinogen, was aspirated with a syringe.

**2.5. Surgical Intervention.** Gingival recessions were treated by CAF associated with a volume-stable collagen matrix (FibroGide, 20 mm height  $\times$  40 mm length  $\times$  6 mm thickness, Geistlich Pharma, Wolhusen, Switzerland) functionalized with i-PRF. Briefly, after local anesthesia with 4% articaine containing 1:100,000 epinephrine, intra-sulcular incisions with a 15C blade (Swann-Morton, Sheffield, England) were performed, involving the premolars and connected to vertical releasing incisions. A split (interproximal papilla)–full (until mucogingival junction)–split (beyond mucogingival junction) thickness flap was elevated (Figure 1(b)). The soft tissue was mobilized with horizontal incisions in the inner part of the flap, and soft tissue perforations accidentally occurred in the distal region of the right canine and left lateral incisor. Root surfaces were mechanically instrumented with periodontal curettes (Hu-Friedy, Chicago, IL, USA) and chemically treated with 24% ethylenediaminetetraacetic acid (EDTA, Straumann PrefGel, Institut Straumann, Basel, Switzerland) for 2 minutes. Chemical preparation of

root surfaces aimed to promote decontamination and improve the blood clot imbrication (adhesion) on the root surface [20, 21]. Interproximal papillae were de-epithelialized. After i-PRF preparation (Intra-Lock L-PRF Intra Spin, Birmingham, AL, USA) [15], a volume-stable collagen matrix was bisected in the long-axis, impregnated with i-PRF for 15 minutes, and positioned in the receptor bed at cemento-enamel junction level (Figures 1(c), 1(d), 1(e), and 1(f)). The tension-free flap was coronally positioned 1 mm beyond the cemento-enamel junction and sutured (5-0 Vicryl; Ethicon, Johnson & Johnson, Somerville, NJ, USA; Figure 1(g)). The matrix was accommodated on the bed, and no sutures were performed.

**2.6. Clinical Outcomes and Follow-Up.** A considerable volume increase was observed at the 5-day follow-up, with no signs of inflammation, dehiscence, or necrosis. Sutures were removed 2 weeks after surgery. The surgical area showed a satisfactory healing 2 months after surgery; however, a volumetric soft tissue reduction was detected in comparison with the immediate post-surgical result (Figure 2(a)). More volumetric decrease was observed at the 6-month follow-up. Nevertheless, the gingival margin remained stable, at the same level from the surgical procedure. At the 9-month re-evaluation, probing depth




FIGURE 3: Baseline and 12-month follow-up measurements. (a) Baseline measurements: root recession area (blue); gingival recession height (black line in the blue area); keratinized tissue band (green); keratinized tissue band height (black line in the green area), (b) 12-month follow-up measurements: root recession area (blue); gingival recession height (black line in the blue area); keratinized tissue band (green); keratinized tissue band height (black line in the green area).

TABLE 1: Measurements of gingival recession area, root coverage area, gingival recession height, root coverage height, and keratinized tissue band height, at baseline and 12 months after the surgical procedure.

|                         | Right canine | Right lateral incisor | Right central incisor | Left central incisor | Left lateral incisor | Left canine |
|-------------------------|--------------|-----------------------|-----------------------|----------------------|----------------------|-------------|
| RRAb (mm <sup>2</sup> ) | 31.13        | 7.96                  | 38.02                 | 28.87                | 11.64                | 32.5        |
| RRAf (mm <sup>2</sup> ) | 3.18         | 0                     | 7.18                  | 8.26                 | 0                    | 13.57       |
| RCa (%)                 | 89.94        | 100                   | 81.11                 | 71.38                | 100                  | 58.24       |
| GRHb (mm)               | 5.02         | 2.09                  | 5.65                  | 3.21                 | 2.6                  | 5.07        |
| GRHf (mm)               | 0.82         | 0                     | 1.73                  | 1.33                 | 0                    | 2.53        |
| RCh (mm)                | 4.2          | 2.09                  | 3.92                  | 1.88                 | 2.6                  | 2.54        |
| KTBHb (mm)              | 6.54         | 6.98                  | 4.89                  | 4.37                 | 5.32                 | 5.85        |
| KTBHf (mm)              | 6.96         | 7.32                  | 6.52                  | 5                    | 4.79                 | 4.92        |
| ΔKTB (mm)               | +0.42        | +0.34                 | +1.63                 | +0.63                | -0.53                | -0.93       |

RRAb, root recession area—baseline; RRAf, root recession area—final; RCa, root coverage area; GRHb, gingival recession height—baseline; GRHf, gingival recession height—final; RCh, root coverage height; KTBHb, keratinized tissue band height—baseline; KTBHf, keratinized tissue band height—final; ΔKTB, keratinized tissue band—difference between initial and final.

was  $\leq 3$  mm in all sites (Figure 2(b)). Gingival margin stability was verified at the 12-month follow-up, with no modification on probing depth, no bleeding on probing, and a stable keratinized mucosa band (Figure 2(c)). Biofilm-induced gingivitis was detected in some teeth at this time. Also, 1 mm of gingival apical migration at the left canine was observed when comparing the 12-month follow-up with the 2-month follow-up. The patient received additional oral hygiene technique instructions, and gingivitis was treated with non-surgical periodontal therapy involving ultrasonic debridement and professional prophylaxis. Differences from baseline to the 12-month follow-up were calculated for the following measurements: gingival recession area, root coverage area, gingival recession height, root coverage height, and keratinized tissue band height (Figure 3, Table 1).

After 30 months of follow-up, clinical parameters were assessed (Figure 4). The probing depths were stable ( $\leq 3$  mm) in comparison with 12-month follow-up; however, bleeding on probing increased, and the patient was diagnosed with biofilm-induced gingivitis. An additional 1 mm recession was verified at the left canine compared to the 12-month follow-up. Oral hygiene instructions were reinforced, and gingivitis was again treated with ultrasonic debridement and professional prophylaxis.

### 3. Discussion

Gingival recessions treated with autogenous connective tissue graft have been associated with postoperative discomfort in the donor area and restrictions in the graft size, which



FIGURE 4: Postoperative follow-up after 30 months. Stable results along the time with minimal modifications. Patient presented bleeding on probing and gingivitis that was further treated during the consultation.

represents relevant limitations for multiple gingival recession treatment [7]. Xenogeneic substitutes have been employed in regenerative periodontal treatment to overcome those limitations. This case report presented the possibility of using a volume-stable collagen matrix functionalized with i-PRF as an alternative for root coverage of multiple RT1 gingival recessions. The proposed approach demonstrated 83% root coverage of multiple gingival recessions in a 12-month follow-up, with a maintenance of the keratinized mucosa band. If we relate our findings with the literature, other substitutes have a lower percentage of root coverage than the present case report [22].

An additional increase in the gingival recession of the left canine was observed 30 months after root coverage. We suppose this rebound over the treatment outcome occurred due to an inflammatory process associated with a gingivitis condition presented by the patient, who missed the follow-up visits for several months [19]. Despite this, our article is the first to show a 30-month follow-up clinical case on CAF associated with a volume-stable collagen matrix functionalized with i-PRF for multiple root coverage treatment. Those findings suggested that the proposed matrix functionalization is a potential strategy for root coverage in multiple gingival recession cases, with a possible reduction in postoperative morbidity, since no access to a donor area for graft collection was necessary.

The functionalized collagen matrix used in this clinical case intends to mimic the natural extracellular matrix of a connective tissue graft, providing a niche for host cells during healing [23, 24], which allows cell migration, adhesion, and differentiation, as well as supporting vascular proliferation. Moreover, it has been suggested that the volume-stable collagen matrix has long-term dimensional stability due to its low level of cross-linking degradation [25]. An *in vitro* study showed that i-PRF induced cell migration and increased the gene expression of TGF- $\beta$  mRNA, platelet-derived growth factor (PDGF), and collagen type I [26]. Theoretically, the PRF may also have an epithelial keratinization capacity [27]. Thus, the use of a volume-stable collagen matrix impregnated with i-PRF could increase the healing potential as compared to autogenous connective tissue graft [28–31].

The success of root coverage depends on the gingival thickness and the amount of keratinized mucosa band. A thick soft tissue provides greater tissue stability and reduces

the recurrence of gingival recession [32]. Studies reported a complete root coverage in cases with an initial gingival thickness  $\leq 1$  mm and a keratinized mucosa band of 2 mm or less when a connective tissue graft was used to augment the soft tissue during root coverage surgery [12, 33]. The present clinical case showed that a satisfactory gingival thickness can be achieved with a functionalized volume-stable collagen matrix with i-PRF, and results were stable for the evaluated period. We speculate that the collagen matrix works as a scaffold for blood clot stabilization, as described in an animal study [34]. The maturation process would result in connective tissue formation with adequate gingival volume. Those findings agree with other studies that prospectively assessed the soft tissue volume increase for long periods [10, 35]. However, our hypothesis requires more studies to be confirmed.

During the follow-up period, a gingival thickness reduction was observed 2 months after the intervention. Matrix thickness was not trimmed before placement in the present clinical case, which may have influenced postoperative results. The large distance between surgical flap and teeth surface may have compromised the vascularization, which could partially explain the incomplete root coverage in some teeth. Another point of consideration was the flap perforations at the right canine and left lateral incisor regions with unpredictable repercussions. Taken together, based on our clinical observations, we suggest that the collagen matrix should be trimmed in its thickness before application on the surgical bed. Other studies also support this recommendation [29, 31].

Considering the limitations of the present case report, it is worth mentioning that, although successful, the proposed approach is limited to one single clinical case. Therefore, our results should be interpreted with caution. Even though a donor site access for connective tissue graft was not necessary in the proposed approach, it is important to highlight that blood collection from the patient in order to produce the PRF is mandatory, which increases morbidity and clinical steps during the surgery. Studies comparing the functionalized collagen matrix with the matrix alone are recommended to draw conclusions on the clinical application of this technique. In addition, this case may have a similar outcome with reduced costs if a CAF alone or in combination with a connective tissue graft was performed [36]. Nevertheless, comparative research would be necessary

to evaluate those aspects. In addition, further randomized clinical studies are mandatory to confirm our findings in larger samples. The molecular events and the underlying mechanisms by which collagen matrices associated with blood derivatives influence tissue formation were not elucidated yet.

#### 4. Conclusion

Based on the findings of this clinical report and considering that results are limited to one single case, the volume-stable collagen matrix functionalized with i-PRF was a feasible approach for multiple RT1 gingival recession treatment, demonstrating 83% of root coverage in a 12-month follow-up period. Stable results were verified for up to 30 months after root coverage. A donor site access for connective tissue harvesting was avoided, which potentially reduced patient morbidity.

#### Conflicts of Interest

The authors declare that they have no conflicts of interest.

#### Acknowledgments

The authors would like to thank Geistlich Pharma do Brasil for the donation of the collagen matrix used in this case report. Roberta Michels and Gabriel Leonardo Magrin received scholarships from the Coordination for the Improvement of Higher Education Personnel (Coordenação para Aperfeiçoamento de Pessoal de Nível Superior—CAPES, Ministry of Education, Brazil) and thanks this support. Roberta Michels also received a scholarship from the Osteology Foundation (Lucerne, Switzerland). Gabriel Leonardo Magrin also received a scholarship from the International Team for Implantology (ITI Foundation, Basel, Switzerland) and a research grant from the Osteology Foundation (Project number: 20-127, Luzern, Switzerland) and would like to thank the financial assistance.

#### References

- [1] P. G. Pini, "Mucogingival deformities," *Annals of Periodontology*, vol. 4, no. 1, pp. 98–101, 1999.
- [2] L. Chambrone and D. N. Tatakis, "Periodontal soft tissue root coverage procedures: a systematic review from the AAP regeneration workshop," *Journal of Periodontology*, vol. 86, no. 2-s, pp. S8–S11, 2015.
- [3] P. Cortellini and N. F. Bissada, "Mucogingival conditions in the natural dentition: narrative review, case definitions, and diagnostic considerations," *Journal of Periodontology*, vol. 89, pp. S204–S213, 2018.
- [4] P. Cortellini and N. F. Bissada, "Mucogingival conditions in the natural dentition: narrative review, case definitions, and diagnostic considerations," *Journal of Clinical Periodontology*, vol. 2018, no. 45, pp. S190–S198, 2016.
- [5] F. Cairo, M. Nieri, S. Cincinelli, J. Mervelt, and U. Pagliaro, "The interproximal clinical attachment level to classify gingival recessions and predict root coverage outcomes: an explorative and reliability study," *Journal of Clinical Periodontology*, vol. 38, no. 7, pp. 661–666, 2011.
- [6] G. P. Pini-Prato, F. Cairo, M. Nieri, D. Franceschi, R. Rotundo, and P. Cortellini, "Coronally advanced flap versus connective tissue graft in the treatment of multiple gingival recessions: a split-mouth study with a 5-year follow-up," *Journal of Clinical Periodontology*, vol. 37, no. 7, pp. 644–650, 2010.
- [7] G. Zucchelli, M. Mele, M. Stefanini et al., "Patient morbidity and root coverage outcome after subepithelial connective tissue and de-epithelialized grafts: a comparative randomized-controlled clinical trial," *Journal of Clinical Periodontology*, vol. 37, no. 8, pp. 728–738, 2010.
- [8] V. Moraschini, M. D. Calasans-Maia, A. T. Dias, F. M. de Carvalho, S. C. Sartoretto, and A. Sculean, "Effectiveness of connective tissue graft substitutes for the treatment of gingival recessions compared with coronally advanced flap: a network meta-analysis," *Clinical Oral Investigations*, vol. 24, no. 10, pp. 3395–3406, 2020.
- [9] F. M. de Carvalho, M. A. Nagasawa, V. Moraschini, J. Ata-Ali, A. Sculean, and J. A. Shibli, "Clinical efficacy of xenogeneic and allogeneic 3D matrix in the management of gingival recession: a systematic review and meta-analysis," *Clinical Oral Investigations*, vol. 24, no. 7, pp. 2229–2245, 2020.
- [10] O. Zuhre, D. Akakpo, P. Eickholz et al., "Tunnel technique with connective tissue graft versus coronally advanced flap with enamel matrix derivate for root coverage: 5-year results of an RCT using 3D digital measurement technology for volumetric comparison of soft tissue changes," *Journal of Clinical Periodontology*, vol. 48, no. 7, pp. 949–961, 2021.
- [11] L. C. Cummings, W. B. Kaldahl, and E. P. Allen, "Histologic evaluation of autogenous connective tissue and acellular dermal matrix grafts in humans," *Journal of Periodontology*, vol. 76, no. 2, pp. 178–186, 2005.
- [12] M. Stefanini, I. Mounssif, S. Barootchi, L. Tavelli, H. L. Wang, and G. Zucchelli, "An exploratory clinical study evaluating safety and performance of a volume-stable collagen matrix with coronally advanced flap for single gingival recession treatment," *Clinical Oral Investigations*, vol. 24, no. 9, pp. 3181–3191, 2020.
- [13] H. M. Jhaveri, M. S. Chavan, G. B. Tomar, V. L. Deshmukh, M. R. Wani, and P. D. Miller, "Acellular dermal matrix seeded with autologous gingival fibroblasts for the treatment of gingival recession: a proof-of-concept study," *Journal of Periodontology*, vol. 81, no. 4, pp. 616–625, 2010.
- [14] F. J. Strauss, J. Nasirzade, Z. Kargarpour, A. Stähli, and R. Gruber, "Effect of platelet-rich fibrin on cell proliferation, migration, differentiation, inflammation, and osteoclastogenesis: a systematic review of in vitro studies," *Clinical Oral Investigations*, vol. 24, no. 2, pp. 569–584, 2020.
- [15] R. J. Miron, M. Fujioka-Kobayashi, M. Hernandez et al., "Injectable platelet rich fibrin (i-PRF): opportunities in regenerative dentistry?," *Clinical Oral Investigations*, vol. 21, no. 8, pp. 2619–2627, 2017.
- [16] R. J. Miron, G. Zucchelli, M. A. Pikos et al., "Use of platelet-rich fibrin in regenerative dentistry: a systematic review," *Clinical Oral Investigations*, vol. 21, no. 6, pp. 1913–1927, 2017.
- [17] R. J. Miron, M. Fujioka-Kobayashi, M. Bishara, Y. Zhang, M. Hernandez, and J. Choukroun, "Platelet-rich fibrin and soft tissue wound healing: a systematic review," *Tissue Engineering. Part B, Reviews*, vol. 23, no. 1, pp. 83–99, 2017.
- [18] J. Choukroun and S. Ghanaati, "Reduction of relative centrifugation force within injectable platelet-rich-fibrin (PRF) concentrates advances patients' own inflammatory cells, platelets and growth factors: the first introduction to the low speed



- centrifugation concept,” *European Journal of Trauma and Emergency Surgery*, vol. 44, no. 1, pp. 87–95, 2018.
- [19] I. L. C. Chapple, B. L. Mealey, T. E. Van Dyke, P. M. Bartold, H. Dommisch, and P. Eickholz, “Periodontal health and gingival diseases and conditions on an intact and a reduced periodontium: consensus report of workgroup 1 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions,” *Journal of Clinical Periodontology*, vol. 45, pp. S68–S77, 2018.
  - [20] J. Blomlöf, L. Jansson, L. Blomlöf, and S. Lindskog, “Root surface etching at neutral pH promotes periodontal healing,” *Journal of Clinical Periodontology*, vol. 23, no. 1, pp. 50–55, 1996.
  - [21] J. Biomiof, L. Biomi, and S. Lindskog, “Effect of different concentrations of EDTA on smear removal and collagen exposure in periodontitis-affected root surfaces,” *Journal of Clinical Periodontology*, vol. 24, no. 8, pp. 534–537, 1997.
  - [22] S. Barootchi, L. Tavelli, G. Zucchelli, W. V. Giannobile, and H. L. Wang, “Gingival phenotype modification therapies on natural teeth: a network meta-analysis,” *Journal of Periodontology*, vol. 91, no. 11, pp. 1386–1399, 2020.
  - [23] L. Ferrantino, D. Bosshardt, M. Nevins, G. Santoro, M. Simion, and D. Kim, “Tissue integration of a volume-stable collagen matrix in an experimental soft tissue augmentation model,” *The International Journal of Periodontics & Restorative Dentistry*, vol. 36, no. 6, pp. 807–815, 2016.
  - [24] D. S. Thoma, R. E. Jung, D. Schneider et al., “Soft tissue volume augmentation by the use of collagen-based matrices: a volumetric analysis,” *Journal of Clinical Periodontology*, vol. 37, no. 7, pp. 659–666, 2010.
  - [25] N. Naenni, P. Walter, C. H. F. Hämmerle, R. E. Jung, and D. S. Thoma, “Augmentation of soft tissue volume at pontic sites: a comparison between a cross-linked and a non-cross-linked collagen matrix,” *Clinical Oral Investigations*, vol. 25, no. 3, pp. 1535–1545, 2021.
  - [26] R. J. Miron, J. Chai, S. Zheng, M. Feng, A. Sculean, and Y. Zhang, “A novel method for evaluating and quantifying cell types in platelet rich fibrin and an introduction to horizontal centrifugation,” *Journal of Biomedical Materials Research. Part A*, vol. 107, no. 10, pp. 2257–2271, 2019.
  - [27] O. Ucak Turer, M. Ozcan, B. Alkaya, S. Surmeli, G. Seydaoglu, and M. C. Haytac, “Clinical evaluation of injectable platelet-rich fibrin with connective tissue graft for the treatment of deep gingival recession defects: a controlled randomized clinical trial,” *Journal of Clinical Periodontology*, vol. 47, no. 1, pp. 72–80, 2020.
  - [28] S. H. Mathes, L. Wohlwend, L. Uebersax et al., “A bioreactor test system to mimic the biological and mechanical environment of oral soft tissues and to evaluate substitutes for connective tissue grafts,” *Biotechnology and Bioengineering*, vol. 107, no. 6, pp. 1029–1039, 2010.
  - [29] D. S. Thoma, C. C. Villar, D. L. Cochran, C. H. F. Hämmerle, and R. E. Jung, “Tissue integration of collagen-based matrices: an experimental study in mice,” *Clinical Oral Implants Research*, vol. 23, no. 12, pp. 1333–1339, 2012.
  - [30] D. S. Thoma, N. Naenni, E. Figuero et al., “Effects of soft tissue augmentation procedures on peri-implant health or disease: a systematic review and meta-analysis,” *Clinical Oral Implants Research*, vol. 29, no. ii, pp. 32–49, 2018.
  - [31] D. S. Thoma, M. Zeltner, M. Hilbe, C. H. F. Hämmerle, J. Hüsler, and R. E. Jung, “Randomized controlled clinical study evaluating effectiveness and safety of a volume-stable collagen matrix compared to autogenous connective tissue grafts for soft tissue augmentation at implant sites,” *Journal of Clinical Periodontology*, vol. 43, no. 10, pp. 874–885, 2016.
  - [32] M. S. Tonetti and S. Jepsen, “Clinical efficacy of periodontal plastic surgery procedures: consensus report of Group 2 of the 10th European Workshop on Periodontology,” *Journal of Clinical Periodontology*, vol. 41, no. Suppl, pp. S36–S43, 2014.
  - [33] M. Stefanini, G. Zucchelli, M. Marzadori, and M. de Sanctis, “Coronally advanced flap with site-specific application of connective tissue graft for the treatment of multiple adjacent gingival recessions: a 3-year follow-up case series,” *The International Journal of Periodontics & Restorative Dentistry*, vol. 38, no. 1, pp. 25–33, 2018.
  - [34] J. C. Imber, A. Roccuzzo, A. Stähli et al., “Immunohistochemical evaluation of periodontal regeneration using a porous collagen scaffold,” *International Journal of Molecular Sciences*, vol. 22, no. 20, p. 10915, 2021.
  - [35] K. Jepsen, M. Stefanini, M. Sanz, G. Zucchelli, and S. Jepsen, “Long-term stability of root coverage by coronally advanced flap procedures,” *Journal of Periodontology*, vol. 88, no. 7, pp. 626–633, 2017.
  - [36] M. Stefanini, K. Jepsen, M. de Sanctis et al., “Patient-reported outcomes and aesthetic evaluation of root coverage procedures: a 12-month follow-up of a randomized controlled clinical trial,” *Journal of Clinical Periodontology*, vol. 43, no. 12, pp. 1132–1141, 2016.